



KWANTLEN POLYTECHNIC UNIVERSITY

CHEMISTRY 1210

Final Exam

April 17, 2014

Time: 3 hours

Name: ANSWERS

Student #: \_\_\_\_\_

**Instructions:**

Answer all questions on the exam paper.

Circle correct answers for multiple choice questions.

Show all work for non-multiple choice problems.

The left hand pages may be used for rough work; these pages will not be graded.

A formula sheet and periodic table are provided; no other materials may be used.


Work independently. Cheating will not be tolerated.

This exam should have 34 questions.



Unless otherwise specified you may assume the temperature is 25°C

Page	Mark	Maximum
2		5
3		6
4		5
5		5
6		6.5
7		4
8		5
9		7.5
10		7
11		6
12		7
13		6
14		8
15		6
16		5
<b>Total</b>		<b>89</b>

1) [5 marks total] The following oxidation-reduction reaction occurs in *basic* solution:




a) [4 marks] Balance the reaction.



b) [1 mark] Which species is the reducing agent?

P<sub>4</sub>

- 2) [3 marks total] A 25.00-mL sample of a solution of  $\text{Mo}^{3+}$  solution was reacted with 0.0600 N  $\text{KMnO}_4$ :



- a) [1 mark] If the  $\text{Mo}^{3+}$  in the reaction came from  $\text{MoCl}_3$  (molar mass 202.3 grams), what is the equivalent mass of the  $\text{MoCl}_3$ ?



$$\frac{202.3 \text{ g}}{\text{mol}} \times \frac{1 \text{ mol}}{3 \text{ equiv}} = \boxed{67.43 \text{ g equiv}}$$

- b) [2 marks] The 25.00 mL sample of  $\text{Mo}^{3+}$  solution required 15.00 mL of the  $\text{KMnO}_4$  solution for complete reaction. Determine the **molarity** of the  $\text{Mo}^{3+}$  in the solution.

$$\frac{15.00 \text{ mL} \times 0.06 \text{ equiv/L} \times \frac{1 \text{ Mo}^{3+}}{3 \text{ equiv}}}{25.00 \text{ mL}} = \boxed{0.012 \text{ M}}$$

- 3) [1 mark] The activation energy for the forward direction of a reaction is 50 kJ and for the reverse direction is 30 kJ. The  $\Delta H$  for the overall reaction is:

- a) -20 kJ  
 b) 20 kJ  
 c) -80 kJ  
 d) 80 kJ  
 e) There is not enough information to answer this question.

- 4) [2 marks] The uncoiling of DNA is a first order process with an activation energy of about 420 kJ/mol. At 50°C the half-life for uncoiling is estimated to be 2 minutes. What is the half-life at a normal body temperature of 37°C?

- a)  $2.8 \times 10^{-3}$  min  
 b) 1.4 min  
 c) 2 min  
 d) 2.85 min  
 e) 1411 min

$$\ln\left(\frac{t_2}{t_1}\right) = \frac{E_a}{R} \left( \frac{1}{T_1} - \frac{1}{T_2} \right)$$

$$\ln\left(\frac{t}{2}\right) = \frac{420,000}{8,314 \text{ J/mol K}} \left( \frac{1}{310.15} - \frac{1}{323.15} \right)$$

$$\ln\left(\frac{t}{2}\right) = 6.55 \Rightarrow t =$$

- 5) [2 marks] A reaction has a rate constant  $k = 1.25 \times 10^{-4} \text{ mol L}^{-1} \text{ s}^{-1}$ . If the initial concentration of the reactant is 0.0750 M, what concentration remains after 10.0 minutes?


- a) 0 M  
b) 0.0696 M  
c) 0.0738 M  
d) 0.0749 M  
e) 0.150 M

$$[A]_t = [A]_0 - kt$$

$$= 0.075 - 1.25 \times 10^{-4} \frac{\text{M}}{\text{s}} \times 600 \text{ s}$$

$$= 0$$

- 6) [3 marks] The rate of the reaction:



was studied using the iodine clock technique and the following data were obtained ( $t$  is the time the reaction took):

Run	$[\text{I}^{-1}]_0 \text{ (M)}$	$[\text{S}_2\text{O}_8^{2-}]_0 \text{ (M)}$	$t \text{ (s)}$
1	0.0400	0.0400	88.0
2	0.0800	0.0400	44.0
3	0.0800	0.0800	22.1

- a) The order of this reaction with respect to  $\text{S}_2\text{O}_8^{2-}$  is:

- i) 0  
ii)  $\frac{1}{2}$   
iii) 1  
iv) 2  
v) 3




- b) The overall order of this reaction is:

- i) 0  
ii)  $\frac{1}{2}$   
iii) 1  
iv) 2  
v) 3

- c) If a run is made with  $[\text{I}^{-1}]_0 = 0.120 \text{ M}$  and  $[\text{S}_2\text{O}_8^{2-}]_0 = 0.0400$ , you would expect the reaction to complete in approximately:

- i) 29 s  
ii) 33 s  
iii) 44 s  
iv) 66 s  
v) 88 s

- 7) [5 marks total] The Raschig reaction produces aqueous hydrazine,  $\text{N}_2\text{H}_4(\text{aq})$ , from  $\text{NH}_3(\text{aq})$  and  $\text{OCl}^-(\text{aq})$  in basic, aqueous solution. A proposed mechanism is:



- a) [3 marks] Determine the rate law based on this reaction mechanism, show all your work clearly.

$$\text{rate} = k_2 [\text{NH}_2\text{Cl}] [\text{NH}_3]$$

$\text{NH}_2\text{Cl}$  is RI, so:

$$k_1 [\text{NH}_3] [\text{OCl}^-] = k_2 [\text{NH}_2\text{Cl}] [\text{NH}_3]$$


$$\Rightarrow [\text{NH}_2\text{Cl}] = \frac{k_1 [\text{OCl}^-]}{k_2}$$

$$\begin{aligned} \text{So rate} &= k_2 [\text{NH}_3] \left( \frac{k_1 [\text{OCl}^-]}{k_2} \right) \\ &= \underline{\underline{k_1 [\text{OCl}^-] [\text{NH}_3]}} \end{aligned}$$

- b) [1 mark] The following substances are intermediates:  $\text{NH}_2\text{Cl}$ ,  $\text{OH}^-$ ,  $\text{N}_2\text{H}_5^+$

- c) [1 mark] This reaction mechanism involves a catalyst (circle one): YES NO

- 8) [2 marks] Find  $K_p$  for the following reaction at 25°C:




$$K_c = 2.4 \times 10^{-3} \text{ at } 25^\circ\text{C}$$

- a)  $3.9 \times 10^{-10}$
- b)  $5.6 \times 10^{-8}$
- c)  $3.9 \times 10^{-6}$
- d)  $5.6 \times 10^{-4}$
- e) 1.5
- f) 100

$$K_p = 2.4 \times 10^{-3} (0.08314 \times 298.15)^{-2} = 3.9 \times 10^{-6}$$

- 9) [2.5 marks total] For the equilibrium mixture at 200°C:





Indicate if the partial pressure of  $\text{NH}_3$  will increase, decrease or remain the same when the following changes are made:


(Circle your choice: I – increase D – decrease NC – no change)

Add some Ne	I	D	NC
Add some $\text{H}_2\text{S}$	I	D	NC
Add some $\text{NH}_4\text{HS}$	I	D	NC
Increase the temperature	I	D	NC
Double the volume of the container (at constant T)	I	D	NC

- 10) [2 marks] Given the following equilibria:



$K_c$  for the equilibrium:



will be:

- a)  $2.5 \times 10^{-24}$
- b)  $5.3 \times 10^{-15}$
- c)  $2.1 \times 10^{-14}$
- d)  $2.3 \times 10^4$
- e)  $4.8 \times 10^{13}$

$$\frac{1.1 \times 10^{-5}}{(4.8 \times 10^{-10})^2} = 4.8 \times 10^{13}$$

11) [2 marks] At a given temperature,  $K_c = 3.24$  for the reaction:



If 0.800 mol of both  $\text{H}_2$  and  $\text{CO}_2$  are placed in a 1.00 L container at this temperature, when the system comes to equilibrium the concentration of  $\text{CO}(\text{g})$  will be:

- a) 1.60 M
- b) 0.800 M
- c) 0.611 M
- ☒ d) 0.514 M
- e) 0.247 M

$$\begin{array}{ccccccc}
 \text{H}_2 & + & \text{CO}_2 & \rightleftharpoons & \text{H}_2\text{O} & + & \text{CO} \\
 0.8 & & 0.8 & & 0 & & 0 \\
 -x & & -x & & +x & & +x \\
 0.8-x & & 0.8-x & & x & & x
 \end{array}$$

$$\frac{x^2}{(0.8-x)^2} = 3.24$$

$$\frac{x}{0.8-x} = 1.8$$

$$x = 1.44 - 1.8x$$

$$2.8x = 1.44$$

$$x = 0.514$$

12) [1 mark] For the reaction:




If a container is filled with these gases, such that  $P(\text{I}_2) = 10.0$  bar,  $P(\text{Br}_2) = 5.0$  bar, and  $P(\text{IBr}) = 20.0$  bar, in which direction will the reaction proceed?

- a) The reaction proceeds to the left.
- ☒ b) The reaction proceeds to the right.
- c) The reaction is at equilibrium.
- d) The reaction volume is required in order to answer this question.
- e) The temperature is required in order to answer this question.

$$Q = \frac{400}{10 \times 5} = 8$$

13) [1 mark] Choose the correct statement when the reaction



is at equilibrium:

- a) The rate constant for the forward reaction is equal to the rate constant for the reverse reaction.
- ☒ b) The rate for the forward reaction is equal to the rate for the reverse reaction.
- c) The concentrations of all reactants and products are equal.
- d) Increasing the volume of the reaction container will increase the yield of ammonia.
- e) The equilibrium expression for the reaction is  $K_c = \frac{[\text{N}_2][\text{H}_2]^3}{[\text{NH}_3]^2}$

14) [2 marks] An indicator ( $pK_a = 5.0$ ) changes colour from yellow to blue. It last appears yellow when  $[Ind^{-1}]/[HInd] = 0.04$  and appears completely blue when  $[Ind^{-1}]/[HInd] = 4$ .

- a) Calculate the pH range over which this indicator changes colour and place those pH values in the appropriate blanks below.

pH range: 3.6 (yellow) to 5.6 (blue)

$$10^{-5} = [H^+](0.04) \quad 10^{-5} = [H^+](4)$$

$$\Rightarrow [H^+] = 2.5 \times 10^{-4} \quad \Rightarrow [H^+]_e = 2.5 \times 10^{-6}$$


$$pH = 3.6 \quad \Rightarrow pH = 5.6$$

- b) Would this be a suitable indicator to use for the titration of  $NH_3$  with  $HCl$ ?  
(Circle your choice.)

Yes

No

15) [1 mark] In the equilibrium system



Bronsted-Lowry theory would designate:

- a)  $NH_3$  and  $H_2O$  as the bases.  
b)  $H_2O$  and  $OH^{1-}$  as a conjugate pair.  
c)  $NH_2^{1-}$  and  $H_3O^{+}$  as the acids.  
d)  $NH_2^{1-}$  and  $H_2O$  as a conjugate pair.  
e)  $NH_3$  as amphiprotic.

16) [2 marks] A 10.0 mL sample of a 0.125 M solution of an unknown monoprotic acid has a  $pH = 2.95$ . What is its ionization constant,  $K_a$ ?

a)  $1.0 \times 10^{-7}$   
b)  $1.3 \times 10^{-6}$   
c)  $1.0 \times 10^{-5}$   
d)  $1.1 \times 10^{-3}$   
e)  $9.8 \times 10^{-3}$

$$HA \rightleftharpoons H^+ + A^-$$

i	0.125	0	0	$\frac{x^2}{0.125-x} = K_a$
c	-x	+x	+x	
e	0.125-x	x	x	


$$x = [H^+]_e = 10^{-2.95}$$

So  $K_a =$



17) [3 marks] Methylamine,  $\text{CH}_3\text{NH}_2$ , has a  $K_b = 3.2 \times 10^{-5}$ . What is its percent ionization in 1.0 and 0.1 M solutions, respectively?

- a) 0.018% and 0.056%  
 b) 0.032% and 0.0032%  
 c) 0.56% and 1.8%  
 d) 0.56% in both  
 e) 0.32% in both




18) [2.5 marks] Match the descriptions given below with one of the mixtures described in questions (a) – (e):

- I – A solution with a pH less than 7 that is not a buffer  
 II – A buffer solution with a pH between 4 and 7  
 III – A solution with a pH of 7  
 IV – A buffer solution with a pH between 7 and 10  
 V – A solution with a pH greater than 7 that is not a buffer

- a) A mixture of 1 mole NaOH and 1 mole NaCl in enough water to make 1.0 L  
 b) A mixture of 1 mole NaCl and 1 mole  $\text{CaCl}_2$  in enough water to make 1.0 L  
 c) A mixture of 1 mole NaF and 0.5 mole HF in enough water to make 1.0 L  
 d) A mixture of 50.0 mL 0.10 M HCl with 25.0 mL 0.10 M  $\text{NH}_3$   
 e) A mixture of 25.0 mL 0.10 M HCl with 50.0 mL 0.10 M  $\text{NH}_3$

V  
 III  
 II  
 I  
 IV

19) [2 marks] What mass of sodium acetate (molar mass 82 g) should be dissolved in 200.0 mL of 0.20 M acetic acid to form a buffer of pH = 5.0?  $K_a$  for acetic acid is  $1.8 \times 10^{-5}$ .



$$b + 1 \times 10^{-5} = 0.19999 (1.8)$$

$$b = 0.359972 \text{ M}$$

$$\frac{x(b+x)}{(0.2-x)} = 1.8 \times 10^{-5}$$

$$x = [\text{H}^+]_e = 1 \times 10^{-5}$$


$$\therefore \frac{(b + 1 \times 10^{-5})}{(0.2 - 1 \times 10^{-5})} = 1.8$$

$$\frac{0.359972 \text{ moles}}{\text{L}} \times 0.2 \text{ L} \times \frac{82 \text{ g}}{\text{mol}} = \boxed{5.9 \text{ g}}$$

- 20) [3 marks] What is the pH at the equivalence point for the titration of 0.10 M benzoic acid by 0.10 M sodium hydroxide? ( $K_a$  for benzoic acid is  $6.3 \times 10^{-5}$ )

$V$  mL of BA will require  $V$  mL of NaOH:

$$[BA] = 0.1 \times \frac{V}{2V} = 0.05M \quad [NaOH] = 0.1 \times \frac{V}{2V} = 0.05M$$




$$\frac{x^2}{0.05} = \frac{1 \times 10^{-14}}{6.3 \times 10^{-5}}$$

$$\Rightarrow x = [OH^-]_e = 2.8 \times 10^{-6}M$$


$$pOH = 5.55$$

$$pH = 8.45$$



- 21) [2 marks] How many moles of  $SrF_2$  will dissolve in 1 L of 0.10 M  $Sr(NO_3)_2$  if  $K_{sp}$  for  $SrF_2$  is  $7.9 \times 10^{-10}$ ?

- a)  $2.8 \times 10^{-5}$
- ☒ b)  $4.4 \times 10^{-5}$
- c)  $7.9 \times 10^{-8}$
- d)  $4.0 \times 10^{-9}$
- e)  $7.9 \times 10^{-9}$



- 22) [2 marks] A solution is 0.120 M in  $Pb^{2+}$ . If the  $K_{sp}$  for  $PbCrO_4 = 1.8 \times 10^{-14}$ . In order to precipitate 99.9% of all the  $Pb^{2+}$  present, the  $[CrO_4^{2-}]$  must be:


- a)  $1.8 \times 10^{-17}$
- b)  $1.8 \times 10^{-14}$
- c)  $1.5 \times 10^{-13}$
- ☒ d)  $1.5 \times 10^{-10}$
- e)  $1.3 \times 10^{-7}$

$$0.120 \times \frac{0.1}{100} = 1.2 \times 10^{-4}$$


$$(1.2 \times 10^{-4})[CrO_4^{2-}]_e = 1.8 \times 10^{-14}$$

$$\Rightarrow [CrO_4^{2-}]_e = 1.5 \times 10^{-10}$$

23) [6 marks total] Given the reaction:



- a) [2 marks] Estimate the bond dissociation energy for the O-O single bond. The bond energy for the  $\text{O}_2$  molecule is 498.3 kJ/mol, and the enthalpies of vaporization of  $\text{H}_2\text{O}_2(\text{aq})$  and  $\text{H}_2\text{O}(\text{l})$  are approximately equal.



- b) [1 mark] Give two reasons why your answer above is an estimate and not an accurate calculation of the bond dissociation energy.

- Bond energies are average values only  
 -  $\Delta H_v$  for  $\text{H}_2\text{O}_2$  and  $\text{H}_2\text{O}$  are not exactly equal.

- c) [3 marks] One litre of a solution initially 0.0100 M in  $\text{H}_2\text{O}_2$  and at 25.000°C is reacted. If all the heat produced in the reaction is retained in the solution, what would be the final temperature? Assume the specific heat capacity of the solution to be 4.184 J g<sup>-1</sup>K<sup>-1</sup>, and that the density of the solution is 1.00 g/mL.

$$0.01 \frac{\text{moles H}_2\text{O}_2}{\text{L}} \times 1 \text{ L} \times \frac{1 \text{ rxn}}{2 \text{ H}_2\text{O}_2} \times \frac{-189.32 \text{ kJ}}{\text{mol rxn}}$$

$$= q_{\text{rxn}} = -946.6 \text{ J}$$


$$q_{\text{rxn}} + q_{\text{soln}} \rightarrow +946.6 \text{ J}$$

$$946.6 \text{ J} = 1000 \text{ g} \times 4.184 \frac{\text{J}}{\text{g}^\circ\text{C}} \times \Delta T$$

$$\Delta T = 0.226^\circ\text{C}$$

$$T_f = 25.226^\circ\text{C}$$

24) [2 marks] Given the reaction:




and that the molar enthalpies of formation of  $\text{CO}_2(\text{g})$  and  $\text{H}_2\text{O}(\text{l})$  are  $-393.5 \text{ kJ}$  and  $-285.8 \text{ kJ}$  respectively, the molar enthalpy of formation of butane ( $\text{C}_4\text{H}_8(\text{l})$ ) is:

- a)  $+2017 \text{ kJ/mol}$
- ☒ b)  $-20.3 \text{ kJ/mol}$
- c)  $-107.6 \text{ kJ/mol}$
- d)  $-2017 \text{ kJ/mol}$
- e)  $+20.3 \text{ kJ/mol}$

$$-2696.9 = 4(-393.5) + 4(-285.8) - x$$

$$\Rightarrow x =$$

25) [8 marks total] For the following system:



$$\Delta H^\circ_{298} = +449.6 \text{ kJ}$$

$$\Delta S^\circ_{298} = +609.62 \text{ J/mol K}$$

a)  $\Delta G^\circ_{298} = 267.9 \text{ kJ}$

i) [2 marks]  $K_{p,298}$  for this reaction is:

- (1)  $1.0 \times 10^{-4696}$
- ☒ (2)  $1.1 \times 10^{-47}$
- (3)  $2.0 \times 10^{-5}$
- (4)  $0.90$
- (5)  $9.1 \times 10^{46}$

$$267900 = -8.3144621 \times 298 \times \ln K_p$$

$$\Rightarrow K_p = 1.1 \times 10^{-47}$$

ii) [2 marks] Determine the value of  $\Delta G_{298}$  when  $P(\text{NH}_3) = 0.0010 \text{ bar}$ ,  $P(\text{H}_2\text{O}) = 0.0020 \text{ bar}$  and  $P(\text{SO}_3) = 0.0020 \text{ bar}$ .

$$Q = (0.001)^2 (0.002)(0.002)$$

$$= 4 \times 10^{-12}$$

$$\Delta G = 267900 + 8.3144621 \times 298 \times \ln(4 \times 10^{-12})$$

$$= 202.9 \text{ kJ}$$

iii) [1 mark] Under the conditions in part (c) the forward reaction is: (circle one)

spontaneous

☒ non-spontaneous

- b) [1 mark] At what temperature will this reaction be at equilibrium under standard conditions?

$$\Delta G^\circ = 0 \Rightarrow \Delta H^\circ - T\Delta S^\circ = 0 \Rightarrow \frac{\Delta H^\circ}{\Delta S^\circ} = T$$

$$T = 737,5 \text{ K}$$

- c) [2 marks]  $K_{p,1000}$  for this reaction will be:

- i)  $4.4 \times 10^{-9}$   
 ii)  $1.0 \times 10^{-14}$   
 iii)  $2.3 \times 10^8$   
 iv)  $9.9 \times 10^{13}$   
 v)  $8.8 \times 10^{835}$

$$\Delta G^\circ = 449600 - 1000(609,62)$$

$$= -160020 \text{ J}$$

$$= -8.3144621 \times 1000 \times \ln K_p$$

$$\Rightarrow K_p = 2.2 \times 10^8$$

- 26) [3 marks total] A concentration cell uses a standard hydrogen electrode (SHE) for one half cell. The other half-cell also uses  $\text{H}_2(\text{g})$  at 1 bar pressure, but the hydrogen ions in solution come from a weak acid. The cell so constructed produces 0.245 V.

- a) [0.5 marks] The SHE is the (circle one): ANODE


CATHODE

- b) [0.5 marks] The process occurring at the SHE is (circle one):

OXIDATION

REDUCTION

- c) [2 marks] Determine the pH of the buffered solution.




$$\text{pH} = 4.14$$

$$0.245 = -\frac{0.059159}{2} \log x^2$$


$$\Rightarrow x = [\text{H}^+] = 7.22 \times 10^{-5}$$

27) [2 marks] Determine the equilibrium constant ( $K_e$ ) for the following reaction at 25°C.




Given the standard reduction potentials:  $\text{Pb}^{2+}/\text{Pb} = -0.125 \text{ V}$  and  $\text{Cu}^{2+}/\text{Cu} = +0.337 \text{ V}$

- a)  $2.41 \times 10^{-16}$
- b)  $6.80 \times 10^{-8}$
- c)  $1.65 \times 10^{-7}$
- d)  $1.46 \times 10^7$
- e)  $4.15 \times 10^{15}$



$$\mathcal{E}^\circ = \frac{+0.059159}{2} \log K \Rightarrow K = 2.4 \times 10^{-16}$$

28) [2 marks] Given the reaction:



What would be the voltage if  $[\text{Ni}^{2+}] = 0.020 \text{ M}$  and  $[\text{Al}^{3+}] = 3.60 \text{ M}$ ?


- a) 1.3880 V
- b) 1.4322 V
- c) 1.4712 V
- d) 1.3488 V
- e) 1.3182 V

$$\begin{aligned} Q &= \frac{3.6^2}{(0.02)^3} = 1.62 \times 10^6 \\ \mathcal{E} &= 1.41 - \frac{0.059159}{6} \log 1.62 \times 10^6 = 1.3488 \text{ V} \end{aligned}$$

29) [2 marks] A copper electrode weighs 35.42 g before electrolysis (of a  $\text{CuSO}_4$  solution) and 36.69 g after. The electrolysis was performed using a current of 3.50 amperes. How long did the electrolysis take?

$$\begin{aligned} \text{a) } 9.2 \text{ s} & \quad (36.69 - 35.42) \times \frac{1 \text{ mol}}{63.546 \text{ g}} \times \frac{2\text{e}^-}{1 \text{ Cu}} \times \frac{96485.3365 \text{ Coul}}{1 \text{ mol}} \\ \text{b) } 275 \text{ s} & \\ \text{c) } 551 \text{ s} & \\ \text{d) } 1102 \text{ s} & \quad \times \frac{1 \text{ s}}{3.5 \text{ Coul}} = 1102 \text{ s} \\ \text{e) } 1.35 \times 10^4 \text{ s} & \end{aligned}$$

30) [2 marks] Given the reaction:



What is  $\Delta G^\circ$ ?

- a) -2108 kJ
- b) -211 kJ
- c) -105 kJ
- d) 211 kJ
- e) 2108 kJ

$$\begin{aligned} \Delta G^\circ &= -20(96485.3365)(1.092) \\ &= -2107 \text{ kJ} \end{aligned}$$

31) [1 mark] The normal boiling point of a liquid:

- a) Is the temperature at which the liquid and vapour are in equilibrium
- b) Varies with the atmospheric pressure
- c) Is the temperature at which the vapour pressure of the liquid is 1 atm
- d) Is the temperature at which the vapour pressure of the liquid equals the external pressure
- e) Is directly proportional to the molar mass of the liquid.

32) [2 marks] At 35°C, the vapour pressure of CS<sub>2</sub> is 512 mmHg, and of acetone, CH<sub>3</sub>COCH<sub>3</sub>, is 344 mmHg. It is known that the acetone-CS<sub>2</sub> intermolecular forces are weaker than the acetone-acetone or CS<sub>2</sub>-CS<sub>2</sub> intermolecular forces. Given this information, you would expect that:

- a) A mixture of 100.0 mL of acetone and 100.0 mL of CS<sub>2</sub> has a volume of 200.0 mL.
- b) A mixture of 100.0 mL of acetone and 100.0 mL of CS<sub>2</sub> has a volume less than 200.0 mL.
- c) When acetone and CS<sub>2</sub> are mixed at 35°C heat is absorbed.
- d) When acetone and CS<sub>2</sub> are mixed at 35°C heat is evolved.
- e) The vapour pressure above the solution would be lower than predicted.

33) [3 marks] The primary constituent of lemon oil is the hydrocarbon, limonene, that is 88.16% C and 11.84% H. A solution of 8.362 g of limonene in 50.00 g of benzene freezes at 2.37°C. Pure benzene freezes at 5.5°C. The freezing point depression constant for benzene is 5.12°C kg/mol. Determine the molecular formula of limonene.

$$\Delta T = 5.5 - 2.37 = 3.13^\circ\text{C}$$

$$3.13 = 1.512 \times m$$

$$\Rightarrow m = 0.6113 \dots$$


$$\text{moles limonene} = 0.6113 \frac{\text{moles}}{\text{kg}} \times 0.05 \text{ kg} = 0.030$$

$$\therefore \text{molar mass limonene} = \frac{8.362 \text{ g}}{0.030 \text{ moles}} = 273.6 \frac{\text{g}}{\text{mol}}$$

$$\left. \begin{array}{l} 273.6 \times 0.8816 \times \frac{1 \text{ mol}}{12.011 \text{ g}} = 20.1 \\ 273.6 \times 0.1184 \times \frac{1 \text{ mol}}{1.0079 \text{ g}} = 32.1 \end{array} \right\} \therefore \text{limonene is } \boxed{\text{C}_{20}\text{H}_{32}}$$

34) [5 marks total] A certain compound has a normal melting point of  $41^{\circ}\text{C}$  and a normal boiling point of  $123^{\circ}\text{C}$ . The triple point is at  $39^{\circ}\text{C}$  and 85 mmHg.

- a) [3 marks] Sketch the phase diagram for this substance. Label all regions, lines and points appropriately.



- b) [1 mark] Does the solid phase of this substance have a density greater or less than that of the liquid phase? How do you know? (No marks for guessing. ☺)

greater - exists at higher P.

- c) [1 mark] Describe what happens when a sample of this substance at 2 atm and a temperature of  $20^{\circ}\text{C}$  is heated at constant pressure to a temperature of  $50^{\circ}\text{C}$ .

Solid warms to about  $42^{\circ}\text{C}$   
 Solid melts at about  $42^{\circ}\text{C}$   
 liquid warms to  $50^{\circ}\text{C}$